

$\varepsilon'_K/\varepsilon_K$ AT NEXT-TO-LEADING IN $1/N_C$ AND TO LOWEST ORDER CHPT

JOHAN BIJNENS

*Department of Theoretical Physics 2, Lund University
Sölvegatan 14A, S 22362 Lund, Sweden*

JOAQUIM PRADES

*Departamento de Física Teórica y del Cosmos, Universidad de Granada
Campus de Fuente Nueva, E-18002 Granada, Spain*

We report on a calculation of $\varepsilon'_K/\varepsilon_K$ at next-to-leading order in the $1/N_c$ expansion and to lowest order in Chiral Perturbation Theory. We also discuss the chiral corrections to our results and give the result of including the two known chiral corrections.

1 Introduction

Recently, direct CP violation in the Kaon system has been unambiguously established by KTeV at Fermilab and by NA48 at CERN¹. The present world average is

$$\text{Re}(\varepsilon'_K/\varepsilon_K) = (19.3 \pm 2.4) \cdot 10^{-4}. \quad (1)$$

Recent reviews and predictions for this quantity in the Standard Model and earlier references are in². Here, we report on a calculation³ of this quantity in the chiral limit and next-to-leading (NLO) order in $1/N_c$. We also discuss the changes when the known chiral corrections -final state interactions (FSI) and $\pi_0 - \eta$ mixing- are included.

Direct CP-violation in the $K \rightarrow \pi\pi$ decay amplitudes is parameterized by

$$\frac{\varepsilon'_K}{\varepsilon_K} = \frac{1}{\sqrt{2}} \left[\frac{A[K_L \rightarrow (\pi\pi)_{I=2}]}{A[K_L \rightarrow (\pi\pi)_{I=0}]} - \frac{A[K_S \rightarrow (\pi\pi)_{I=2}]}{A[K_S \rightarrow (\pi\pi)_{I=0}]} \right]. \quad (2)$$

$K \rightarrow \pi\pi$ amplitudes can be decomposed into definite isospin amplitudes as

$$\begin{aligned} i A[K^0 \rightarrow \pi^0 \pi^0] &\equiv \frac{a_0}{\sqrt{3}} e^{i\delta_0} - \sqrt{\frac{2}{3}} a_2 e^{i\delta_2}, \\ i A[K^0 \rightarrow \pi^+ \pi^-] &\equiv \frac{a_0}{\sqrt{3}} e^{i\delta_0} + \frac{a_2}{\sqrt{6}} e^{i\delta_2}. \end{aligned} \quad (3)$$

with δ_0 and δ_2 the FSI phases.

We want to predict a_0 and a_2 to NLO order in $1/N_c$ and to lowest order in CHPT.

2 Short-Distance Scheme and Scale Dependence

The procedure to obtain the Standard Model effective action $\Gamma_{\Delta S=1}$ below the W -boson mass has become standard and explicit calculations have been performed to two-loops⁴. The full process implies choices of short-distance scheme, regulators, and operator basis. Of course, physical matrix elements cannot depend on these choices.

The Standard Model $\Gamma_{\Delta S=1}$ effective action at scales ν somewhat below the charm quark mass, takes the form⁵

$$\Gamma_{\Delta S=1} \sim \sum_{i=1}^{10} C_i(\nu) \int d^4x Q_i(x) + \text{h.c.} \quad (4)$$

where $Q_i(x)$ are four-quark operators and $C_i = z_i + \tau y_i$ are Wilson coefficients. In the presence of CP-violation, $\tau \equiv -V_{td}V_{ts}^*/V_{ud}V_{us}^*$ gets an imaginary part.

At low energies, it is more convenient to use an effective action $\Gamma_{\Delta S=1}^{LD}$ which uses different degrees of freedom. Different regulators and/or operator basis can be more practical too. The effective action $\Gamma_{\Delta S=1}^{LD}$ depends on all these choices and in particular on the scale μ_c introduced to regulate the divergences, analogous to ν in (4) and on effective couplings g_i , which are the equivalent of the Wilson coefficients in (4). Matching conditions between the effective field theories

of (4) and $\Gamma_{\Delta S=1}^{LD}$ are obtained by requiring that S -matrix elements of asymptotic states are the same at some perturbative scale.

$$\langle 2|\Gamma_{\Delta S=1}^{LD}|1\rangle = \langle 2|\Gamma_{\Delta S=1}|1\rangle. \quad (5)$$

The matching conditions fix *analytically* the short-distance behavior of the couplings g_i

$$g_i(\mu_c, \dots) = \mathcal{F}(C_i(\nu), \alpha_s(\nu), \dots). \quad (6)$$

This was done explicitly in ⁶ for $\Delta S = 2$ transitions and used in ³ for $\Delta S = 1$ transitions.

2.1 The Heavy X -Boson Method

For energies below the charm quark mass, we use an effective field theory of heavy color-singlet X -bosons coupled to QCD currents and densities^{3,6,7}. For instance, the effective action reproducing

$$Q_1(x) = [\bar{s}\gamma^\mu(1 - \gamma_5)d] [\bar{u}\gamma_\mu(1 - \gamma_5)u](x)$$

is

$$\Gamma_X \equiv g_1(\mu_c, \dots) \int d^4y X_1^\mu \{ [\bar{s}\gamma_\mu(1 - \gamma_5)d](x) + [\bar{u}\gamma_\mu(1 - \gamma_5)u](x) \}. \quad (7)$$

Here the degrees of freedom of quarks and gluons above the scale μ_c have been integrated out. The advantage of this method is that two-quark currents are unambiguously identified and that QCD densities are much easier to match than four-quark operators.

We use a 4-dimensional Euclidean cut-off μ_c to regulate UV divergences. We can now calculate $\Delta S = 1$ Green's functions with the X -boson effective theory consistently.

3 Long-Distance–Short-Distance Matching

Let's study the $\Delta S = 1$ two-point function

$$\Pi(q^2) \equiv i \int d^4x e^{iq \cdot x} \langle 0 | T(P_i^\dagger(0) P_j(x) e^{i\Gamma_X} | 0 \rangle.$$

The P_i are pseudoscalar sources with quantum numbers describing $K \rightarrow \pi$ amplitudes.

Taylor expanding the off-shell amplitudes $K \rightarrow \pi$ obtained from these Green's

functions, in external momentum and π , and K masses, one can obtain the couplings of the CHPT Lagrangian. These predict $K \rightarrow \pi\pi$ at a given order. This is unambiguous.

At leading order in the $1/N_c$ expansion the contribution to the Green function $\Pi(q^2)$ is factorizable. This only involves strong two-point functions and is model independent.

The non-factorizable contribution, is NLO in the $1/N_c$ expansion. It involves the integration ⁸ of strong four-point functions $\Pi_{P_i P_j J_a J_b}$ over the momentum Euclidean r_E that flows through the currents/densities J_a and J_b from 0 to ∞ , schematically written as

$$\Pi(q^2) \sim \int \frac{d^4 r_E}{(2\pi)^4} \Pi_{P_i P_j J_a J_b}(q_E, r_E). \quad (8)$$

We separate long- from short-distance physics with a cut-off μ in r_E . The short distance part can be treated within OPE QCD.

Recently, it was emphasized that dimension eight operators may be numerically important for low values of the cut-off scale⁹. This issue can be studied straightforwardly in our approach.

There is *no* model dependence in our evaluation of $K \rightarrow \pi$ amplitudes at NLO in $1/N_c$ within QCD up to now. The long distance part from 0 up to μ remains. For very small values of μ one can use CHPT but it starts to be insufficient already at relatively small values of μ . Too small to match with the short-distance part. The first step to enlarge the CHPT domain is to use a good hadronic model for intermediate energies. We used the ENJL model¹⁰. It has several good features -it includes CHPT to order p^4 , for instance- and also some drawbacks as explained in ⁶. Work is in progress to implement the large N_c constraints on three- and four-point functions along the lines of ¹¹.

4 ε'_K in the Chiral Limit

To a very good approximation,

$$|\varepsilon'_K| \simeq \frac{1}{\sqrt{2}} \frac{\text{Re } a_2}{\text{Re } a_0} \left\{ -\frac{\text{Im } a_0}{\text{Re } a_0} + \frac{\text{Im } a_2}{\text{Re } a_2} \right\}. \quad (9)$$




Figure 1. Plot showing the matching between short- and long-distances for $\varepsilon'_K/\varepsilon_K$. Notice the wide plateau. Curves I and II correspond to different choices in the running of α_s to two-loops.

The lowest order CHPT values for $\text{Re } a_0$ and $\text{Re } a_2$ are obtained from a fit¹² to $K \rightarrow \pi\pi$ and $K \rightarrow \pi\pi\pi$ amplitudes to order p^4 . Our results⁷ reproduce the $\Delta I = 1/2$ enhancement within 40 %. We use the *experimental* lowest order CHPT values¹² for $\text{Re } a_I$ to predict ε'_K as shown in Figure 1.

For the two dominant operators and $\varepsilon'_K/\varepsilon_K$ we obtain at NLO in $1/N_c$ and in the chiral limit^{3,7}

$$\begin{aligned} B_{6\chi}^{(1/2)NDR}(2\text{ GeV}) &= 2.5 \pm 0.4 \\ B_{8\chi}^{(3/2)NDR}(2\text{ GeV}) &= 1.35 \pm 0.20 \\ \left| \frac{\varepsilon'_K}{\varepsilon_K} \right|_{\chi} &= (60 \pm 30) \cdot 10^{-4}. \end{aligned} \quad (10)$$

5 Higher Order CHPT Corrections

The rôle of FSI in the standard² predictions of $\varepsilon'_K/\varepsilon_K$ has been recently studied¹³. We took a different strategy. The ratio $\text{Im } a_I/\text{Re } a_I$ has no FSI to all orders, thus FSI only affects the ratio $\text{Re } a_2/\text{Re } a_0$.

Among the isospin breaking effects only π^0 - η mixing is under control and is known¹⁴ to order p^4 . Other real p^4 and higher electromagnetic corrections are mostly unknown.

Including π^0 - η mixing and FSI our result (10) becomes

$$\left| \frac{\varepsilon'_K}{\varepsilon_K} \right| = (34 \pm 18) \cdot 10^{-4}. \quad (11)$$

Acknowledgments

This work is partially supported by the European Union TMR Network EURODAPHNE (Contract No. ERBFMX-CT98-0169), the Swedish Science Foundation (NFR), CICYT, Spain (Grant No. AEN-96/1672), and Junta de Andalucía (Grant No. FQM-101).

References

1. A. Alavi-Harati et al., *Phys. Rev. Lett.* **83**, 22 (1999); V. Fanti et al., *Phys. Lett. B* **465**, 335 (1999); A. Ceccucci, CERN Particle Physics Seminar (29 February 2000) <http://www.cern.ch/NA48>
2. A.J. Buras et al., hep-ph/0007313; M. Ciuchini and G. Martinelli, hep-ph/0006056; S. Bertolini et al, hep-ph/0002234; T. Hambye et al., *Nucl. Phys. B* **564**, 391 (2000); E. Pallante et al, València preprint IFIC/00-31
3. J. Bijnens and J. Prades, *J. High Energy Phys.* **06**, 035 (2000)
4. A.J. Buras et al, *Nucl. Phys. B* **400**, 37, 75 (1993); M. Ciuchini et al, *Nucl. Phys. B* **415**, 403 (1994)
5. A.J. Buras, hep-ph/9806471.
6. J. Bijnens and J. Prades, *J. High Energy Phys.* **01**, 002 (2000)
7. J. Bijnens and J. Prades, *J. High Energy Phys.* **01**, 023 (1999)
8. J. Bijnens and J. Prades, *Phys. Lett. B* **342**, 331 (1995); *Nucl. Phys. B* **444** 523 (1995)
9. V. Cirigliano et al, hep-ph/0007196
10. J. Bijnens, *Phys. Rep.* **265**, 369 (1996) and references therein
11. S. Peris, M. Perrottet, and E. de Rafael, *J. High Energy Phys.* **05**, 011 (1998)
12. J. Kambor, J. Missimer, and D. Wyler, *Phys. Lett. B* **261**, 496 (1991)
13. E. Pallante and A. Pich, *Phys. Rev. Lett.* **84**, 2568 (2000); hep-ph/0007208
14. G. Ecker et al, *Phys. Lett. B* **477**, 88 (2000)